




# Evolution of thermal physiology alters the projected range of threespine stickleback under climate change

Sara J. Smith<sup>1,2,3</sup>  | Stephanie Mogensen<sup>1</sup> | Tegan N. Barry<sup>1</sup> | Antoine Paccard<sup>4</sup>  | Heather A. Jamniczky<sup>5</sup> | Rowan D. H. Barrett<sup>4</sup> | Sean M. Rogers<sup>1,6</sup> 

<sup>1</sup>Department of Biological Sciences, University of Calgary, Calgary, Alberta, Canada

<sup>2</sup>Informatics Group, Harvard University, Cambridge, Massachusetts, USA

<sup>3</sup>Department of Biology, Boston University, Boston, Massachusetts, USA

<sup>4</sup>Redpath Museum and Department of Biology, McGill University, Montreal, QC, Canada

<sup>5</sup>Department of Cell Biology & Anatomy, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>6</sup>Bamfield Marine Sciences Centre, Bamfield, British Columbia, Canada

## Correspondence

Sara J. Smith, Biology, Mount Royal University, 4825 Mt Royal Gate SW, Calgary, AB T3E 6K6, Canada.  
Email: [ssmith6@mtroyal.ca](mailto:ssmith6@mtroyal.ca)

## Present address

Sara J. Smith, Biology, Mount Royal University, Calgary, Canada

## Funding information

Alberta Innovates Technology Futures; Natural Sciences and Engineering Research Council of Canada; Canada Research Chairs

Handling Editor: Andrew Kinziger

## Abstract

Species distribution models (SDMs) are widely used to predict range shifts but could be unreliable under climate change scenarios because they do not account for evolution. The thermal physiology of a species is a key determinant of its range and thus incorporating thermal trait evolution into SDMs might be expected to alter projected ranges. We identified a genetic basis for physiological and behavioural traits that evolve in response to temperature change in natural populations of threespine stickleback (*Gasterosteus aculeatus*). Using these data, we created geographical range projections using a mechanistic niche area approach under two climate change scenarios. Under both scenarios, trait data were either static (“no evolution” models), allowed to evolve at observed evolutionary rates (“evolution” models) or allowed to evolve at a rate of evolution scaled by the trait variance that is explained by quantitative trait loci (QTL; “scaled evolution” models). We show that incorporating these traits and their evolution substantially altered the projected ranges for a widespread panmictic marine population, with over 7-fold increases in area under climate change projections when traits are allowed to evolve. Evolution-informed SDMs should improve the precision of forecasting range dynamics under climate change, and aid in their application to management and the protection of biodiversity.

## KEYWORDS

climate change, critical thermal limits, evolution, species distribution models, thermal biology, threespine stickleback

## 1 | INTRODUCTION

Temperature is a powerful driver of global biogeography, and species distributions frequently reflect temperature gradients in both aquatic and terrestrial habitats (Hochachka & Somero, 2002). Many species adopt thermal strategies (such as thermoregulation or acclimation) that determine their thermal niche (Coutant, 1987; Huey & Kingsolver, 1989; Huey & Slatkin, 1976) and thermal traits can

provide a target for directional selection if the environment changes to include temperatures outside the range encompassed by the thermal niche. Adaptation can thus permit species to persist at temperatures that would have previously led to extirpation (Hoffman & Sgrò, 2011; Sexton et al., 2009). Under moderate climate change scenarios, mean global oceanic temperature is projected to increase in excess of 2°C by the end of the century (IPCC, 2014), with more extreme changes predicted in localized regions (Eyer et al., 2019;

Walther et al., 2002) for both cooling and warming events (Hu et al., 2018). Accurately predicting species range patterns under climate change therefore requires data for temperature-associated adaptive trait evolution (Bay et al., 2017; Bush et al., 2016; Catullo et al., 2015).

The genetic basis of thermal traits has been explored in many terrestrial ectotherms (Angilletta et al., 2002; Bowler & Terblanche, 2008; Dillon et al., 2009; Rolandi et al., 2018) and there have been integrative frameworks proposed for how best to use this information to assess the vulnerability of organisms to environmental changes (Araújo et al., 2013; Bay et al., 2017; Huey et al., 2012). Given the differences in thermal biology between terrestrial and aquatic ectotherms (Sunday et al., 2011) and differential warming between terrestrial and aquatic habitats (IPCC, 2014, 2018), predicting how the thermal biology of aquatic organisms will respond to environmental changes requires an understanding of the genetic architecture underlying ecologically relevant thermal traits (Healy et al., 2018) in aquatic systems. The genetic basis of thermal traits has been explored in a number of socioeconomically and culturally important fish species (Everett & Seeb, 2014; Jackson et al., 1998; Jin et al., 2017; Larson et al., 2016; Muñoz et al., 2014; Perry et al., 2001) and there has been an extensive body of work on the thermal biology of the common killifish (*Fundulus heteroclitus*; Bryant et al., 2018; Fanguie et al., 2006, 2009; Healy et al., 2018). These studies suggest that a significant proportion of phenotypic variance in thermal traits can be explained by genetic variation (e.g., 35 single nucleotide polymorphisms [SNPs] explained 51.9% of the variation for upper thermal tolerance in killifish; Healy et al., 2018), highlighting the potential for heritable responses to changes in thermal conditions.

While the climate and ocean temperatures are warming overall, thermal events characterized by both cold and heat extremes are occurring with increasing frequency (Hu et al., 2018; IPCC, 2014, 2018; Stott, 2016). Regionally downscaled environmental changes can be highly spatially heterogeneous relative to global temperature trends, and are likely to be the most relevant for species-specific responses in the context of contemporary climate change (Walther et al., 2002). For example, range expansions of marine species have been mediated by both ocean-scale warming and regional cooling (Zeidberg & Robison, 2007). In such cases, population persistence and adaptation will rely on both upper and lower critical thermal limits, forcing organisms to respond independently to extreme heat and extreme cold in separate events (Herring et al., 2015, 2018, 2020; Hu et al., 2018; Walther et al., 2002). Importantly, adaptation in one thermal trait can shift the thresholds of correlated thermal traits (Buckley & Huey, 2016; Denny & Dowd, 2012; Hoffman & Sgrò, 2011; Huey & Kingsolver, 2011). Associations between traits influence the rate of trait adaptation and thus correlated traits must be considered when attempting to predict future species distributions under climate change (Bestion et al., 2015; Hoffman & Sgrò, 2011; Valladares et al., 2014). Recently there have been steps to incorporate theoretical trait evolution into species distribution models (SDMs) (e.g., by including the breeder's equation for a key phenotypic trait; Bush et al.,

2016; Catullo et al., 2015; Kearney & Porter, 2009). However, no model has used empirical estimates of evolutionary rates to inform projected species ranges under climate change scenarios.


Threespine stickleback (*Gasterosteus aculeatus*, Figure 1a) is a useful vertebrate species for understanding the impact of adaptation on range dynamics under climate change. This species exhibits its widespread phenotypic variation (Hendry et al., 2013), there are publicly available genomic resources (Jones, Grabherr, et al., 2012) and prior research on temperature-associated evolutionary rates for this species (Barrett et al., 2011; Morris et al., 2018). It is also feasible to artificially breed multiple hybrid generations in common garden laboratory environments. Moreover, there is a regionally downscaled climate model for the Pacific Northwest (Alexander et al., 2018; IPCC, 2014; Reynolds et al., 2007), which makes panmictic marine populations of *G. aculeatus* in the eastern Pacific Ocean (Kirch et al., 2021; Morris et al., 2018) an ideal system for testing how adaptive trait variation could affect projections of species ranges under climate change.

The objective of this study was to incorporate ecologically and evolutionarily relevant data on thermal traits into end-of-century projections of SDMs. To accomplish this, we assessed physiological and behavioural temperature-associated traits in two marine and two freshwater populations of threespine stickleback, as well as within-population  $F_1$  families and marine-freshwater hybrid  $F_1$  and  $F_2$  families. We used the hybrid families to construct genetic linkage maps and identified multiple quantitative trait loci (QTL) associated with thermal preference, as well as upper and lower critical thermal limits. Using these genetically based traits and empirically derived evolutionary rate data for this species (Barrett et al., 2011), as well as others (Morgan et al., 2020; Sanderson et al., 2021), we constructed SDMs using a mechanistic niche area approach that integrated the trait data reported in this study and allowed those traits to evolve under two climate change scenarios, three distinct types of evolutionary models and three evolutionary rates.

## 2 | MATERIALS AND METHODS

### 2.1 | Sample collection and husbandry

We collected adult *Gasterosteus aculeatus* from two marine populations (Bamfield, M1, 48°49'12.69" N, 125° 8'57.90" W; Garden Bay Lagoon, M2, 49°37'52.84" N, 124° 1'49.26" W) and two freshwater populations (Hotel Lake, FW1, 49°38'26.94" N, 124° 3'0.69" W; Klein Lake, FW2, 49°43'32.47" N, 123°58'7.83" W) in southwestern British Columbia (Figure 1b) at depths of less than 0.5 m. To ensure that any genetic variation identified would be relevant for adaptation within this region, we confined our study to a single genetically panmictic cluster of populations (Morris et al., 2018). Individuals were maintained in a flow-through system and under a photoperiod that mimicked the natural source population environment during collection periods before transport. We transported



**FIGURE 1** (a) Adult threespine stickleback (*Gasterosteus aculeatus*) from a single genetic cluster were sampled from (b) two marine and two freshwater populations in the Canadian Pacific Northwest. These populations were assayed for (c) thermal preference (green), critical thermal minima (CTmin, blue) and maxima (CTmax, red). Thermal trait values for marine populations (M1 and M2) were incorporated into the species distribution models [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/mec.16396)]

the fish to the Life and Environmental Sciences Animal Resources Centre at the University of Calgary, where we separated the fish into population-specific 113-L glass aquaria at a density of ~20 fish per aquarium. We acclimated marine individuals to freshwater salinity over 1 week and maintained fish in a common garden environment (salinity of 4–6 ppt, water temperature of  $15 \pm 2^\circ\text{C}$ , and a photoperiod of 16:8 h light:dark). Individuals were allowed to acclimate for at least 2 weeks before experiments (1 week for stress reduction post-transfer, 1 week for common garden environment acclimation and salinity ramp). Each common garden aquarium was on a closed system with individual filters, air stones and water supply. We fed all adult fish *ad libitum* once per day with thawed bloodworms (Hikari Bio-Pure Frozen Bloodworms). All collections and transfers were approved by Fisheries and Oceans Canada (marine collections and provincial transfers), the Ministry of Forests, Lands, and Natural Resource Operations (freshwater collections) and the Huu-ay-aht First Nations (marine collections).

## 2.2 | Crossing design for marine and freshwater $F_1$ families

We collected eggs from females and fertilized the eggs with extracted testes from euthanized males. We transferred the fertilized egg mass to a mesh-bottomed egg incubator suspended in a 37-L aquarium for hatching. Each hatching aquarium was maintained with a single air stone and a filter. Once hatched, we reared the larval fish in 37-L hatching aquaria until they reached a total length (TL) of ~1 cm, after which we split the families into family-specific 113-L aquaria to maintain suitable densities. We fed the larval fish *ad libitum* twice daily with live *Artemia* spp. nauplii, and then gradually transitioned the diet to chopped, thawed bloodworms (Hikari Bio-Pure Frozen Bloodworms) *ad libitum* once daily as they reached ~2 cm TL. The  $F_1$  families were maintained in a common garden environment identical to that of the  $F_0$  populations. We produced one  $F_1$  family for each population (M1\_ $F_1$ , M2\_ $F_1$ , FW1\_ $F_1$ , and FW2\_ $F_1$ ).

## 2.3 | Crossing design for hybrid mapping families

To generate genetically heterogeneous marine–freshwater  $F_1$  families from wild  $F_0$  parents, we collected eggs from marine females and fertilized the eggs with extracted testes from euthanized freshwater males. Egg masses were hatched and juveniles were reared as detailed above. We produced one  $F_1$  family of  $M1 \times FW1$  hybrids (hereafter referred to as  $H1_{F_1}$ ) and three  $F_1$  families of  $M1 \times FW2$  hybrids (hereafter referred to as  $H2_{F_1}$ ). The hybrid  $F_1$  families were maintained in a common garden environment identical to that of the  $F_0$  populations. To generate  $F_2$  families for linkage map construction and QTL mapping, we crossed siblings from the same  $F_1$  family using the same methodology that was used to generate the  $F_1$  families. The amount of trait variation explained by these hybrid crosses is likely to be higher than would be observed in marine–marine or freshwater–freshwater crosses, increasing our likelihood of detecting QTL. Overall, we produced one  $F_2$  family of  $H1 \times H1$  hybrids (referred to as  $H1_{F_2}$ ) and three families of  $H2 \times H2$  hybrids (referred to as  $H2_{F_2-1}$ ,  $H2_{F_2-2}$  and  $H2_{F_2-3}$ ). All  $F_2$  individuals were raised as described above in a common garden environment identical to that of the  $F_0$  and  $F_1$  individuals to ensure consistent thermal history.

## 2.4 | Thermal tolerance and preference experiments

To assess the lower and upper limits of physiological thermal tolerance, we conducted standard critical thermal minimum (CTmin) and maximum (CTmax) experiments on adult fish (Barrett et al., 2011; Fangué et al., 2006; Hutchison, 1961). At these sublethal limits, the fish experience a loss of equilibrium (LOE) at which they lose the ability to escape conditions that would ultimately lead to their death (Beitinger et al., 2000). Before each experiment, individuals were fasted for 24 h. Our experimental tank held 1000-ml glass beakers aerated individually to prevent thermal stratification. After a 15-min acclimation to the experimental apparatus in the individual beakers, we cooled or heated the water (for CTmin or CTmax, respectively) at a rate of  $\sim 0.33^\circ\text{C min}^{-1}$ . We assessed wild  $F_0$  individuals ( $n_{M1} = 32$ ,  $n_{M2} = 14$ ,  $n_{FW1} = 15$ ,  $n_{FW2} = 16$ ,  $N = 77$ ; Figure S2), and laboratory raised  $F_1$  ( $n_{M1_{F_1}} = 13$ ,  $n_{M2_{F_1}} = 15$ ,  $n_{FW1_{F_1}} = 15$ ,  $n_{FW2_{F_1}} = 15$ ,  $N = 58$ ; Figure S2), and  $F_2$  individuals ( $n_{H1_{F_2}} = 28$ ,  $n_{H2_{F_2-1}} = 36$ ,  $n_{H2_{F_2-2}} = 21$ ,  $n_{H2_{F_2-3}} = 17$ ,  $N = 102$ ; Figure S3). All individuals were assessed for CTmin, allowed to recover for at least 3 days, and then assessed for CTmax to keep thermal stress history consistent. The onset of erratic behaviours associated with a behavioural stress response (i.e., “agitation windows”; Turko et al., 2020) occurred below  $5.0^\circ\text{C}$  and above  $25.0^\circ\text{C}$  during CTmin and CTmax experiments, respectively. Normal behaviour was observed between  $5.0$  and  $25.0^\circ\text{C}$ , whereas outside of those temperatures, individuals gradually exhibited more extreme stress responses (e.g., increased gilling

rate, erratic movement, muscle spasms, listing; as outlined by the Canadian Council of Animal Care guidelines) until reaching LOE and the inability of an individual to right itself (the experimental endpoint, measured in  $0.5^\circ\text{C}$  increments; Barrett et al., 2011; Fangué et al., 2006; Hutchison, 1961).

To assess the range of behavioural thermoregulation, we conducted thermal preference experiments in a temperature Shuttlebox (Loligo Systems). Experimental pools were set with a static gradient of  $10^\circ\text{C}$  in the cool side and  $20^\circ\text{C}$  in the warm side, connected with a water bridge at  $15^\circ\text{C}$  (acclimation temperature). Temperature was continually monitored and recorded by immersed temperature probes, connected to a computer-driven temperature controller and data acquisition system (DAQ-M; Loligo Systems). Movement in the Shuttlebox was tracked by an infrared-sensitive uEye USB 2.0 camera (IDS Imaging Development Systems) which connected to SHUTTLESOFT (version 2; Loligo Systems) and allowed monitoring of fish movement. Following recovery from CTmax experiments (at least 3 days), individuals were acclimated in the Shuttlebox for 15 min, with entry at “cool” or “warm” pool randomized, to allow for exploration of the gradient. After acclimation to the experimental apparatus, the movement of each individual was tracked in 1-s intervals for 30 min, recording the preferred ambient water temperature (thermal preference). Individuals were assessed individually in the Shuttlebox apparatus to avoid any confounding social behaviours (e.g., as seen in shoaling behaviours; Cooper et al., 2018). At the time of data collection for thermal trait experiments, all individuals were adults.


Relationships between all thermal traits separated by population and generation were assessed using the Pearson product-moment correlation with the *corrplot* (version 0.84, Wei et al., 2017) and *Hmisc* (version 3.14-0, Harrell, 2014) packages in R (R Core Team, 2021).

## 2.5 | Isolation and characterization of SNPs

Genomic DNA was extracted from caudal fin tissue using a phenol-chloroform-based protocol. We digested tissues overnight in digestion buffer and proteinase K at  $55^\circ\text{C}$ , then performed multiple phenol–chloroform and ethanol washes to isolate the DNA. We assessed the quantity of the extracted DNA using the Quant-iT PicoGreen dsDNA assay kit (ThermoFisher Scientific) and Synergy HT plate reader with the Gen5-associated software (BioTek). We prepared restriction site-associated DNA (RAD) libraries (Peterson et al., 2012) using *MluCI* and *NlaIII* restriction enzymes (New England Biolabs), ligation of individual barcodes and pooling of 48 individuals per library at equimolar concentrations. We performed a final polymerase chain reaction to amplify DNA and add library-specific indices to allow for pooling of multiple libraries. We sequenced three libraries at the McGill University and Génome Québec Innovation Center on one lane of an Illumina HiSeq 4000 (Illumina).

**TABLE 1** Significant quantitative trait loci (QTL) for critical thermal minimum (CTmin), critical thermal maximum (CTmax) and thermal preference (Preference) traits with the percent of trait variance explained (PVE) for each QTL, the logarithm of odds (LOD) threshold for significance and the LOD peak observed for each QTL

QTL Name	Family	Phenotype	Linkage group	Position (cM)	95% CI	PVE (%)	LOD threshold (5%)	LOD peak
Q1	H1_F2	CTmax	XII	4.56	0.000–11.38	45.2	2.71	2.87
Q2	H1_F2	Preference	III	50.04	43.21–70.80	53.3	2.78	3.64
Q3	H2_F2_1	CTmin	III	11.57	6.347–52.62	28.9	2.55	2.63
Q4	H2_F2_2	CTmin	XXI	55.67	37.01–66.87	53.7	2.75	3.01
Q5	H2_F2_2	CTmax	XX	81.36	73.12–88.48	83.3	5.57	7
Q6	H2_F2_2	Preference	I	108.2	83.54–109.2	69.3	3.27	4.62
Q7	H2_F2_3	Preference	VII	31.48	24.02–53.87	87.1	3.61	5.78



**FIGURE 2** Linkage maps constructed for quantitative trait loci (QTL) analyses for (a) H1 F<sub>2</sub> hybrids (M1 × FW1) and (b) H2 F<sub>2</sub> hybrids (M1 × FW2) families, rendered with LINKAGEMAPVIEW (Ouellette et al., 2018), with locations of QTL indicated (Table 1) [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.com)]

## 2.6 | Assembly of the genetic linkage maps

After barcode demultiplexing and filtering out low-quality reads in STACKS (Catchen et al., 2013), we removed PCR duplicates from the raw sequences and aligned these cleaned sequences to the *G. aculeatus* reference genome (Jones, Grabherr, et al., 2012) using the Burrows–Wheeler transform (Li & Durbin, 2010). Individual libraries were concatenated and filtered (Puritz et al., 2014) using *vcftools* (version 3.0, Danecek et al., 2011) and split into chromosome-specific variant call format (VCF) files using *SnpSift* (Cingolani et al., 2012) to assemble the linkage maps chromosome by chromosome. We assigned markers to a linkage group with an initial logarithm of the odds (LOD) score of 3 after filtering out markers that showed high levels of segregation distortion and missing observations (>20% missing data) in LEP-MAP3 (Rastas, 2017). Unassigned markers were subsequently added to the existing linkage group at a LOD score of 3 and a minimum size limit of five markers per linkage group. We ordered the markers using a minimum posterior value of 0.001 and collapsed multiple markers when the probability difference between markers was <0.01 (Rastas, 2017). The final linkage map was phased in LEP-MAP3 (Rastas, 2017) and subset for use in R (R Core Team, 2021) to generate a list of informative SNPs to use in subsequent analyses with the *qtl* package (version 1.44-9, Broman et al., 2003). The final linkage maps were similar across families in length and spacing between markers, though the H2\_F<sub>2</sub> map had a higher density of markers (Table S1).

## 2.7 | Quantitative trait loci mapping

We analysed the four F<sub>2</sub> families separately with the same methodology to assess the presence of QTL associated with the thermal traits and avoid the confounding effects of unique meiotic events. We calculated conditional genotype probabilities using a hidden Markov model, allowing for possible genotyping errors at a level of 0.0001 using a Kosambi mapping function with a fixed step width, prior to running genome scans with a single QTL model (Arends et al., 2014; Broman & Sen, 2009). We determined the LOD score significance thresholds for each trait through permutation tests for each family (5000 permutations per chromosome). We pulled significant QTL above the genome-wide significance threshold ( $\alpha = 0.05$ ; Greenwood et al., 2011), calculated confidence intervals of QTL location based on nearby markers, and estimated the percent variance explained by each QTL peak marker ( $PVE = 1 - 10^{(-2 \cdot LOD)/n}$ ). We identified two QTL associated with CT<sub>min</sub>, two QTL associated with CT<sub>max</sub> and three QTL associated with thermal preference (Table 1, Figure 2).

## 2.8 | Environmental variables used to construct SDMs

We compiled environmental data widely used in the construction of SDMs to estimate suitable habitat in both present-day and

end-of-century forecasts (Wiens et al., 2009), including bathymetry, sea ice extent and concentration, salinity, and sea surface temperature (SST). We used 2014 data as our baseline year to match the forecasting baseline of the Fifth Assessment Report (IPCC, 2014). We assumed a suitable habitat range for this species in the Pacific Northwest to consist of coastal areas (water depth <200 m) where sea ice is never present (i.e., no sea ice at the maximum extent). *G. aculeatus* has a broad salinity tolerance (Bayly, 2003; Divino et al., 2016) and salinity is not limiting in any part of the range (Zweng et al., 2013), so salinity was not included in the final present-day or forecasted models. We obtained bathymetry data from the General Bathymetric Chart of the Oceans (GEBCO) of the British Oceanographic Data Centre (Weatherall et al., 2015), and maximum sea ice extent data from the Multisensory Analyzed Sea Ice Extent – Northern Hemisphere (MASIE-NH) product (Fetterer et al., 2010). We obtained maximum and minimum daily statistical mean SST from Reynolds et al. (2007) to create rasters of the mean upper quartile, lower quartile, absolute minimum, absolute maximum and median temperatures for the baseline year (2014). We found the minimum, maximum and median temperature for each 1-km<sup>2</sup> grid cell across the range, then reduced the area based on the constraints of sea ice extent and bathymetry. Stickleback thermal trait data were used to set the limits of the ranges within the possible area delineated by sea ice-free water of a suitable depth. Thermal trait measurements were based on our experimental findings reported in the present study.

In the end-of-century forecast for suitable habitat, we assumed bathymetry to be consistent with the modern scenario. In contrast, because the Arctic Ocean is projected to be predominantly free of sea ice in the summer by the end of the century (Johannessen et al., 2004), with significant end-of-century reductions in winter/spring sea ice concentration (reduced to a concentration of 0.1 at the Seward Peninsula; Johannessen et al., 2004), we conservatively set the maximum northern extent of the suitable habitat to the western tip of the Seward Peninsula (65°35'N). The water temperatures were increased based on projections for large marine ecosystems of Northern Oceans from global climate models (Alexander et al., 2018; IPCC, 2014). Raster and maps were created in R (version 4.1.2, R Core Team, 2021) using the packages *raster* (version 3.5.15, Hijmans et al., 2020) and *rgeos* (version 0.5.9, Roger et al., 2020).

## 2.9 | Trait inclusion in SDMs

We incorporated experimental data on the wild marine populations (Figure 1c) to understand how inclusion of trait data may affect range projections under climate change. These trait-defined areas were overlain on the suitable habitat background to delineate projected presence based on thermal traits in both current-day and IPCC-projected Representative Concentration Pathways (RCPs) 4.5 and 8.5. The trait values were kept constant from current day to end-of-century (i.e., not changed) in the “no evolution” projections. In contrast, in the “evolution” projections, we allowed traits

to change based on empirically derived evolutionary rate estimates. We allowed CT<sub>min</sub> to evolve at a rate of 0.63 *haldanes* estimated from a selection experiment previously conducted on populations belonging to the same admixed genetic cluster as the sampled populations (Barrett et al., 2011; Morris et al., 2018). At present, there are no empirical estimates of CT<sub>max</sub> evolution for threespine stickleback. Thus, as a proxy we used a known estimate of CT<sub>max</sub> evolution in fish, which was observed in zebrafish (0.19 *haldanes*; Morgan et al., 2020). Given that the evolutionary rates for CT<sub>min</sub> and CT<sub>max</sub> from Barrett et al. (2011) and Morgan et al. (2020) are quite high relative to other recorded evolutionary rates for phenotypic traits (particularly physiological traits; Sanderson et al., 2021), we also explored the influence of evolution on projected ranges when evolution occurred at more “typical” rates. To do so, we allowed both CT<sub>min</sub> and CT<sub>max</sub> to evolve at a mean rate from a large meta-analysis of phenotypic trait evolution (0.14 *haldanes*; Sanderson et al., 2021).

Since the rates of evolution used here are estimates based on whole-organism tolerance (Barrett et al., 2011; Morgan et al., 2020), and trait evolution is constrained by the underlying genetic architecture of the trait (Barghi et al., 2020; Fisher, 1930; Orr, 1998; Ungerer & Riesebero, 2003), we next explored how selection acting on discrete, identifiable loci could impact evolutionary rates, and the subsequent effects on projected habitat range. To do so, we constructed models that scaled the evolutionary rate of CT<sub>min</sub> or CT<sub>max</sub> based on the observed genetic architecture from our QTL mapping (i.e., we scaled by the percent trait variation that is explained by each locus; “scaled evolution” projections). However, an important restriction in the evolution of CT<sub>min</sub> for all models incorporating trait evolution (“evolution” and “scaled evolution” projections) was a hard boundary at 0°C, with the assumption that population persistence in a sub-zero environment would require additional adaptations to CT<sub>min</sub> improvement (e.g., extreme adaptations observed in Antarctic notothenioid fishes; Cheng & Detrich, 2007; Detrich et al., 2000; Shin et al., 2014). Similarly, we defined a hard boundary for CT<sub>max</sub> evolution at 42°C (Morgan et al., 2018), with the assumption that the strength of selection for CT<sub>max</sub> at this rate would result in decreased phenotypic variation and acclimatory ability leading to an upper thermal limit plateau (Gilchrist & Huey, 1999; Hangartner & Hoffmann, 2016; Morgan et al., 2020).

To quantify the differences in estimated suitable habitat under current-day and end-of-century conditions, we compared areas for each warming scenario to the equivalent scenario under current conditions. Similarly, to compare the differences in evolutionary scenarios, the area of each end-of-century evolutionary trajectory was compared to either the contrasting RCP projection or scaled evolution projection. For these comparisons, we used a North Pole Lambert azimuthal equal area projection for all maps, and georeferenced to known landmarks in ARCGIS version 10.8 (Environmental Systems Research Institute, 2017) to calculate area from the maps generated in R (version 4.1.2, R Core Team, 2021; conversion ratio of 7873.42).

## 3 | RESULTS

### 3.1 | Genetic basis of thermal traits


Threespine stickleback exhibited a wide thermal tolerance range bounded by a mean CT<sub>min</sub> of 1.89°C ( $\pm 1.04^\circ\text{C}$  SD) and a mean CT<sub>max</sub> of 30.1°C ( $\pm 1.78^\circ\text{C}$  SD) (Figure 1c; Figure S2), with these physiological traits being highly correlated ( $r = .79$ , Figure S1). These marine populations also tolerated a wide range of temperatures within which there was no observable stress response (5.0–25.0°C). Thermal preference values were slightly above the acclimation temperature, but with more variance around the mean than is observed in the physiological traits ( $16.7 \pm 3.19^\circ\text{C}$ , Figure 1c; Figure S2). To determine if these measured thermal traits have a genetic basis and could therefore evolve in response to natural selection, we raised hybrid marine-freshwater F<sub>1</sub> ( $N = 2$ ) and F<sub>2</sub> ( $N = 4$ ; Figure S3) families under common garden conditions and used these fish for genome-wide linkage map construction (Table S1) and QTL mapping. Using 41,840 high-quality single nucleotide variants generated from RAD sequencing, we identified two significant QTL for CT<sub>min</sub>, two significant QTL for CT<sub>max</sub> and three significant QTL for thermal preference (Table 1, Figure 2). The trait variance that each QTL explained ranged from 28.9% (CT<sub>min</sub> QTL on linkage group III) to 87.1% (thermal preference QTL on linkage group VII).

### 3.2 | Geographical range of projected suitable habitat

Based on sea ice extent and bathymetry alone, our present-day suitable habitat model suggests a marine range for this population from the southern Bering Sea to northern Washington State, and along the southeast Alaskan Panhandle (combined shaded area in Figure 3a, ~654,122 km<sup>2</sup>). End-of-century IPCC projections resulted in a substantial increase in the overall suitable habitat area for stickleback, with an 873,059-km<sup>2</sup> increase in total area (combined shaded area in Figure 3b–e, ~1,527,181 km<sup>2</sup>) in association with a reduction in sea ice concentration at the northern end of the range.

### 3.3 | “No evolution” SDM

We used our observed values of CT<sub>min</sub>, CT<sub>max</sub> and thermal preference to first inform the boundaries of three distinct environmental regions in the SDMs: (i) a “Normal Behaviour” (NB) area that included environmental temperatures associated with an absence of an observable behavioural stress response (5.0–25.0°C), (ii) a “Within Physiological Limits” (WPL) area that included environmental temperatures within the range of their measured physiological limits (1.8–30.1°C) and (iii) an “Outside Physiological Limits” (OPL) area with environmental temperatures that fall outside the measured physiological limits (below 1.8 and above



**FIGURE 3** Changes in the projected range of marine threespine stickleback (*Gasterosteus aculeatus*) in the northeast Pacific Ocean as a result of incorporating thermal traits into models of (a) current-day environmental conditions, and under IPCC end-of-century projections RCP 4.5 and 8.5, either without trait evolution (b and c; “no evolution” model, orange box) and with trait evolution at a rate of 0.63 *haldanes* for critical thermal minimum (CTmin) and a rate of 0.19 *haldanes* for critical thermal maximum (CTmax) (d and e; “evolution” model, red box). Sampling locations are in the southeast area of the range presented here [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

30.1°C). When we include thermal trait data from the wild marine populations, nearly the entire range of suitable habitat was unaffected by thermal tolerance limits, apart from a slight restriction at the northern end of the range (Figure 3a). However, when restricted to the NB area, the range becomes confined to the west of the northern tip of Kodiak Island (Figure 3a), a limit coinciding with the northern-most known marine population in the Pacific Northwest genetic cluster (Morris et al., 2018). When temperature increases as projected by RCP 4.5, the “no evolution” model projects a 1,011,949-km<sup>2</sup> increase in the NB area within this newly suitable habitat (Figure 3b) when compared to the NB area in the current-day model. Under RCP 8.5, the entirety of suitable habitat area remains within tolerable limits in the “no evolution” model (Figure 3c), with a smaller proportion of the range (10.7%) falling outside of the NB area as compared to the NB area under RCP 4.5 conditions.

### 3.4 | “Evolution” SDMs

Incorporating the evolution of CTmin and CTmax into the SDMs (“evolution” models) results in a large increase in the proportion of suitable habitat that falls within the NB area. At the rate of 0.63 *haldanes* for CTmin (Barrett et al., 2011) and 0.19 *haldanes* for CTmax, the entire range of the suitable habitat range falls within the NB area (Figure 3d,e). These represent a 311,974- and 155,680-km<sup>2</sup> increase in the NB area relative to the “no evolution” models under RCP 4.5 and RCP 8.5, respectively, and an increase in 1,271,032 km<sup>2</sup> in the NB area when compared to the current-day model. If we only allow one of each of these traits to evolve at a time, we find that only CTmin evolution results in a change to range distributions compared to “no evolution” models (Figure S4). Reducing the rate of evolution for these traits to the mean rate of trait evolution reported in Sanderson et al. (2021) still results in all suitable habitat falling



within the NB area under both RCP 4.5 and RCP 8.5. Similarly, scaling CT<sub>min</sub> to evolve at a rate equivalent to the response expected if selection acted solely on the detected CT<sub>min</sub> QTL on either linkage group III (PVE = 28.9%) or XXI (PVE = 53.7%) resulted in identical range distributions as the full “evolution” models. Given that CT<sub>max</sub> evolution had no impact on these range projections, we did not construct “scaled evolution” models for this trait.

## 4 | DISCUSSION

In this study, we have produced the first “evolution-informed” SDMs that incorporate empirical data for temperature-associated traits and their evolution to understand how end-of-century projections change with the inclusion of these data. We assessed physiological (critical thermal minimum, [CT<sub>min</sub>] and maximum [CT<sub>max</sub>]), and behavioural (thermal preference) thermal traits for threespine stickleback from wild marine and freshwater populations, as well as hybrid F<sub>1</sub> and F<sub>2</sub> families. We provide novel characterization of QTL underlying these traits in this species, which are likely to be important genetic targets of selection under climate change. Finally, we incorporated these trait data and three empirical estimates of evolutionary rates into mechanistic niche area SDMs under two climate change scenarios. When trait data are included in projections but held constant (i.e., “no evolution” models), there is an ~6-fold (RCP 4.5) and 7-fold (RCP 8.5) increase in the geographical area of the NB area. The geographical ranges projected for the end-of-century increased by over 7-fold when CT<sub>min</sub> was allowed to evolve at observed trait-specific rates in the “evolution” models. Additionally, when a slower rate of CT<sub>min</sub> was used or the rate was scaled by the percent of the trait variance that is explained by individual loci (reflecting a scenario where selection acted solely on those loci), this substantial increase in NB area was still attained by the end of the century under either RCP scenario. These changes to the projected species ranges under climate change underline the importance of incorporating behavioural as well as physiological data into SDMs (Sunday et al., 2012), as well as the key role that thermal trait evolution could play in range shifts (Buckley et al., 2010; Evans et al., 2015; Lyon et al., 2019).

Our models project a northward range expansion under climate change. This is unsurprising because climate change opens newly available thermal niche space in waters north of the current-day geographical range (Alexander et al., 2018) (as is seen in the “no evolution” model northward expansion, Figure 3b,c). Northward range expansion with climate change due to increasing habitat availability has also been documented in birds (Melles et al., 2011; Rushing et al., 2020; Tingley et al., 2009; Tombre et al., 2019), plants (D'Andrea et al., 2009), other fishes (Fosshiem et al., 2015; Spies et al., 2020; Yapıcı et al., 2016) and pest species (e.g., ticks and mountain pine beetle; Clow et al., 2017; Kurz et al., 2008; Ogden et al., 2006; Sagurova et al., 2019; Sambaraju et al., 2019), as well as in large-scale analyses of diverse taxa assessing the “fingerprints” of climate

change impacts (Parmesan & Yohe, 2003; Platts et al., 2019). What is perhaps less expected is that our models reveal that evolution of cold tolerance, but not heat tolerance, had a substantial impact on predicted ranges under climate change, despite end-of-century climate change scenarios projecting an overall warmer, not cooler, world. This result occurs because even with climate change, waters within the marine range of this stickleback population are not projected to reach temperatures that would surpass their current CT<sub>max</sub> limit. However, there is reason to think that the relevance of cold tolerance in climate change scenarios is not specific to this system and could be generally important to consider. Climate change is leading to an increase in the frequency of extreme temperature events (IPCC, 2014; Stott, 2016), including both extreme heat and extreme cold (Herring et al., 2015, 2018, 2020), which could drive selection on both CT<sub>min</sub> and CT<sub>max</sub> (Buckley & Huey, 2016; Denny & Dowd, 2012; Hoffman & Sgrò, 2011; Kingsolver et al., 2011). Moreover, although northern waters are warming on average, there has been considerable spatial heterogeneity in temperature change (Walther et al., 2002). Importantly, northern waters are not predicted to be uniformly warmer under climate change (Alexander et al., 2018; IPCC, 2014) relative to the southern waters in this range, especially when seasonal variation is considered (Alexander et al., 2018). Seasonal variation and more extreme thermal events under climate change scenarios may force organisms to respond independently to both extreme heat and extreme cold in separate events (Herring et al., 2015, 2018, 2020; Walther et al., 2002), and coupled with the correlation between CT<sub>min</sub> and CT<sub>max</sub> (as observed here, Figure S1), population persistence will rely on adaptation in both upper and lower critical limits. Further investigations to test the empirical rate of evolution of thermal behaviour, physiology and the molecular underpinnings of these key traits would be well served by assessing additional subpopulations along the latitudinal gradient inhabited by stickleback to gain a more detailed understanding of these temperature-associated traits over a wider environmental range.

In interpreting the projections we present here, it is important to note that using a different methodology to assess thermal limits could result in different thresholds relating to tolerance and behaviour (Moyano et al., 2017). In addition, in the absence of an empirical estimate of the rate of evolutionary change for CT<sub>max</sub> in threespine stickleback, we are using a rate observed for this trait in zebrafish as a proxy, which will necessarily be inaccurate for stickleback. It is unclear whether the true rate of CT<sub>max</sub> evolution in stickleback might be slower or more rapid than the rate for zebrafish. For instance, it is possible that in threespine stickleback CT<sub>max</sub> could evolve at similar rates as observed for CT<sub>min</sub>, particularly given similarities in genetic architecture and the correlation between these traits in this study. Although incorporating evolution of CT<sub>max</sub> had no impact on our projected ranges because temperatures under RCP 4.5 and RCP 8.5 never exceeded current CT<sub>max</sub> values, we suspect this trait will be of major importance in many other systems where populations reside in environments closer to their current upper thermal limits (e.g., Deutsch et al.,

2008; Morgan et al., 2019; Nati et al., 2021). Ultimately, given the dramatic temperature changes that are projected for end-of-century marine systems (IPCC, 2014, 2018), there is a critical need for empirical measures of evolutionary rates in thermal traits for predicting future species ranges.

While this set of mechanistic niche models builds upon current SDM work by incorporating functional traits critical to population persistence as well as varied evolutionary scenarios based on empirical measurements, there are limitations to these projections. We present niche models of geographical species ranges, but future work in this area would benefit from including a robust metric of species distribution and population genetic structure within these mechanistic geographical ranges. Deriving distributional maps from statistical algorithms that rely on species occurrence and bioclimatic variables would move beyond binary outcomes of habitat suitability and provide a map of suitable habitat that is scored as an index and could underlie the static and evolving trait areas. Species distributions that build upon these evolution-informed species ranges could be an important tool for projecting population persistence under climate change.

The efficiency of translating the selection acting on a trait into evolutionary responses across generations will depend on the genetic architecture of the trait and the extent of environmental change (Dittmar et al., 2016; Orr, 2005; Rogers et al., 2012). Among-trait genetic correlations can also either speed or slow adaptive evolution depending on whether correlations are reinforcing or antagonistic to the direction of selection (Etterson & Shaw, 2001). In this case, we observe a positive correlation between CTmin and CTmax that could facilitate selection for increased temperature tolerance at both upper and lower limits. However, the empirical rates of trait evolution that we incorporated into our SDMs were estimated using change in phenotypic variation across generations (Barrett et al., 2011; Morgan et al., 2020), rather than evolution at underlying loci. As such, it is likely that some proportion of the observed phenotypic change was due to plastic responses. In our “scaled evolution” models, we took a conservative approach by restricting phenotypic evolution to occur only through heritable change via the loci shown here to be associated with the trait. The large-effect loci that we identified are consistent with expectations from theory that suggest prolonged bouts of adaptation with gene flow (as expected in this system; Jones, Chan, et al., 2012; Rogers et al., 2012; Schluter & Conte, 2009; Schluter et al., 2010, 2021) will favour architectures characterized by fewer, larger effect, more tightly linked alleles (Griswold & Whitlock, 2003; Via et al., 2012; Yeaman & Otto, 2011; Yeaman & Whitlock, 2011). However, the effects of the QTL identified from these populations are probably overestimated, while other loci may have gone undetected (*sensu* the Beavis Effect; Beavis, 1994). These loci may have been detected in some  $F_2$  families and not in others because each mapping family has its own specific meiotic events that enable the identification of certain loci. Additionally, there are many key functional traits that are relevant to population persistence under a changing climate, and it is possible that the traits focused on

here, or unexplored traits related to thermal performance, could display antagonistic pleiotropy.

Collectively, the inclusion of thermal traits and their evolution alters the projected ranges of marine threespine stickleback, with a substantial increase in the projected area that the species may occupy under climate change forecasts. Many traits are evolving in response to climate change (Eliason et al., 2011; Gómez-Ruiz & Lacher, 2019; Horton et al., 2020; Hovel et al., 2016) and SDMs that do not take trait data (and trait evolution) into account could provide less accurate predictions about future species distributions under climate change (Bush et al., 2016; Guisan et al., 2013; Huey et al., 2012; Kearney & Porter, 2004)—an issue of particular concern for species at risk and pest species undergoing range expansion (Bebber, 2015; Cullingham et al., 2011; McLeod et al., 2012). Our results provide a framework for addressing this problem, which will have critical implications for the application of these models in policy, sustainable resource management and the protection of biodiversity in a changing climate.

## ACKNOWLEDGMENTS

The species collections took place on Huvu-ay-aht and Sechelt First Nations traditional territories and we are grateful to them for the opportunity to conduct research in protected and sacred areas. We thank the Bamfield Marine Sciences Centre for the resources required to conduct this research, Sam Owens for the photo in Figure 1a, Peter Peller for creating Figure 1b, Carolyn Tepolt for providing the high-resolution SST data set from Reynolds et al. (2007) during a government shutdown, as well as Kiyoko Gotanda for calculating and providing mean haldane values from the Sanderson et al. (2021) database. We thank Daniel Wuitchik, Sam Yeaman, Jennifer Sunday and Patrick Nosil for feedback on the manuscript, and the anonymous peer reviewers for improving this work. This research was funded by Discovery Grants from the Natural Sciences and Engineering Research Council of Canada to RDHB and SMR, a Canada Research Chair to RDHB, and a New Faculty Award from Alberta Innovates Technology Futures to SMR.

## AUTHOR CONTRIBUTIONS

This study was designed by Sara J. Smith, Rowan D. H. Barrett and Sean M. Rogers; fish husbandry and breeding by Sara J. Smith and Tegan N. Barry; experimental data collection by Sara J. Smith; DNA sequencing and initial processing by Antoine Paccard; bioinformatic and QTL analyses by Sara J. Smith; species distribution modelling by Sara J. Smith and Stephanie Mogensen; the manuscript was written by Sara J. Smith, Rowan D. H. Barrett and Sean M. Rogers, with input from all authors; the study was funded by Heather A. Jamniczky, Rowan D. H. Barrett and Sean M. Rogers.

## OPEN RESEARCH BADGES



This article has earned an Open Data Badge for making publicly available the digitally-shareable data necessary to reproduce the reported

results. The data is available at <https://doi.org/10.5061/dryad.70rxwdbz6> and [https://github.com/sjswuitchik/gasAcu\\_qtl\\_sdm](https://github.com/sjswuitchik/gasAcu_qtl_sdm)

#### DATA AVAILABILITY STATEMENT

The data associated with this work are available as follows: ddRAD DNA sequences: available on NCBI (BioProject [PRJNA795470](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA795470)). Experimental data associated with thermal trait data: available on Dryad (<https://doi.org/10.5061/dryad.70rxwdbz6>). Annotated code and example files: available on GitHub ([https://github.com/sjswuitchik/gasAcu\\_qtl\\_sdm](https://github.com/sjswuitchik/gasAcu_qtl_sdm)).

#### ORCID

Sara J. Smith  <https://orcid.org/0000-0002-1913-0396>

Antoine Paccard  <https://orcid.org/0000-0003-0738-5540>

Sean M. Rogers  <https://orcid.org/0000-0003-0851-8050>

#### REFERENCES

- Alexander, M. A., Scott, J. D., Friedland, K. D., Mills, K. E., Nye, J. A., Pershing, A. J., & Thomas, A. C. (2018). Projected sea surface temperatures over the 21st century: Changes in the mean, variability and extremes for large marine ecosystem regions of Northern Oceans. *Elementa*, 6. <https://doi.org/10.1525/elementa.191>
- Angilletta, M. J., Niewiarowski, P. H., & Navas, C. A. (2002). The evolution of thermal physiology in ectotherms. *Journal of Thermal Biology*, 27, 249–268. <https://doi.org/10.2741/E148>
- Araújo, M. B., Ferri-Yáñez, F., Bozinovic, F., Marquet, P. A., Valladares, F., & Chown, S. L. (2013). Heat freezes niche evolution. *Ecology Letters*, 16(9), 1206–1219. <https://doi.org/10.1111/ele.12155>
- Arends, D., Prins, P., Broman, K. W., & Jansen, R. C. (2014). Tutorial-multiple-QTL mapping (MQM) analysis for R/qtl. <http://www.rqtl.org/tutorials/MQM-tour.pdf>
- Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: A unifying framework to understand positive selection. *Nature Reviews Genetics*, 21(12), 769–781. <https://doi.org/10.1038/s41576-020-0250-z>
- Barrett, R., Paccard, A., Healy, T., Bergek, S., Schulte, P., Schluter, D., & Rogers, S. (2011). Rapid evolution of cold tolerance in stickleback. *Proceedings of Royal Society B*, 278(1703), 233–238. <https://doi.org/10.1098/rspb.2010.0923>
- Bay, R. A., Rose, N., Barrett, R., Bernatchez, L., Ghalambor, C. K., Lasky, J. R., Brem, R. B., Palumbi, S. R., & Ralph, P. (2017). Predicting responses to contemporary environmental change using evolutionary response architectures. *The American Naturalist*, 189(5), 463–473. <https://doi.org/10.1086/691233>
- Bayly, I. A. E. (2003). Salinity tolerance and osmotic behavior of animals in athalassic saline and marine hypersaline waters. *Annual Review of Ecology and Systematics*, 3(1), 233–268. <https://doi.org/10.1146/annurev.es.03.110172.001313>
- Beavis, W. (1994). The power and deceit of QTL experiments: Lessons from comparative QTL studies. *49th Annual Corn & Sorghum Research Conference*, pp. 250–266.
- Bebber, D. P. (2015). Range-expanding pests and pathogens in a warming world. *Annual Review of Phytopathology*, 53, 335–356. <https://doi.org/10.1146/annurev-phyto-080614-120207>
- Beitinger, T., Bennett, W., & McCauley, R. (2000). Temperature tolerances of North American freshwater fishes exposed to dynamic changes in temperature. *Environmental Biology of Fishes*, 58, 237–275. <https://doi.org/10.1023/A:1007676325825>
- Bestion, E., Clobert, J., & Cote, J. (2015). Dispersal response to climate change: Scaling down to intraspecific variation. *Ecology Letters*, 18(11), 1226–1233. <https://doi.org/10.1111/ele.12502>
- Bowler, K., & Terblanche, J. S. (2008). Insect thermal tolerance: What is the role of ontogeny, ageing and senescence? *Biological Reviews*, 83, 339–355. <https://doi.org/10.1111/j.1469-185X.2008.00046.x>
- Broman, K. W., & Sen, S. (2009). *A guide to QTL mapping with R/qtl* (Vol. 46). Springer.
- Broman, K. W., Wu, H., Sen, S., & Churchill, G. A. (2003). R/qtl: QTL mapping in experimental crosses. *Bioinformatics*, 19(7), 889–890. <https://doi.org/10.1093/bioinformatics/btg112>
- Bryant, H. J., Chung, D. J., & Schulte, P. M. (2018). Subspecies differences in thermal acclimation of mitochondrial function and the role of uncoupling proteins in killifish. *The Journal of Experimental Biology*. <https://doi.org/10.1242/jeb.186320>
- Buckley, L., & Huey, R. (2016). How extreme temperatures impact organisms and the evolution of their thermal tolerance. *Integrative and Comparative Biology*, 56(1), 98–109. <https://doi.org/10.1093/icb/icw004>
- Buckley, L., Urban, M., Angilletta, M., Crozier, L., Rissler, L., & Sears, M. W. (2010). Can mechanism inform species' distribution models? *Ecology Letters*, 13, 1041–1054. <https://doi.org/10.1111/j.1461-0248.2010.01479.x>
- Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., McEvey, S., & Ferrier, S. (2016). Incorporating evolutionary adaptation in species distribution modelling reduces projected vulnerability to climate change. *Ecology Letters*, 19(12), 1468–1478. <https://doi.org/10.1111/ele.12696>
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An analysis tool set for population genomics. *Molecular Ecology*, 22(11), 3124–3140. <https://doi.org/10.1111/mec.12354>
- Catullo, R. A., Ferrier, S., & Hoffmann, A. A. (2015). Extending spatial modelling of climate change responses beyond the realized niche: Estimating, and accommodating, physiological limits and adaptive evolution. *Global Ecology and Biogeography*, 24(10), 1192–1202. <https://doi.org/10.1111/geb.12344>
- Cheng, C. H. C., & Detrich, H. W. (2007). Molecular ecophysiology of Antarctic notothenioid fishes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1488), 2215–2232. <https://doi.org/10.1098/rstb.2006.1946>
- Cingolani, P., Patel, V. M., Coon, M., Nguyen, T., Land, S. J., Ruden, D. M., & Lu, X. (2012). Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a new program, SnpSift. *Frontiers in Genetics*, 3(1). <https://doi.org/10.3389/fgen.2012.00035>
- Clow, K. M., Leighton, P. A., Ogden, N. H., Lindsay, L. R., Michel, P., Pearl, D. L., & Jardine, C. M. (2017). Northward range expansion of *Ixodes scapularis* evident over a short timescale in Ontario, Canada. *PLoS One*, 12(12), 1–15. <https://doi.org/10.1371/journal.pone.0189393>
- Cooper, B., Adriaenssens, B., & Killen, S. S. (2018). Individual variation in the compromise between social group membership and exposure to preferred temperatures. *Proceedings of the Royal Society B: Biological Sciences*, 285(1880). <https://doi.org/10.1098/rspb.2018.0884>
- Coutant, C. C. (1987). Thermal preference: When does an asset become a liability? *Environmental Biology of Fishes*, 18(5), 161–172. <https://doi.org/10.1007/BF00000356>
- Cullingham, C. I., Cooke, J. E. K., Dang, S., Davis, C. S., Cooke, B. J., & Coltman, D. W. (2011). Mountain pine beetle host-range expansion threatens the boreal forest. *Molecular Ecology*, 20(10), 2157–2171. <https://doi.org/10.1111/j.1365-294X.2011.05086.x>
- D'Andrea, L., Broennimann, O., Kozłowski, G., Guisan, A., Morin, X., Keller-Senften, J., & Felber, F. (2009). Climate change, anthropogenic disturbance and the northward range expansion of *Lactuca serriola* (Asteraceae). *Journal of Biogeography*, 36(8), 1573–1587. <https://doi.org/10.1111/j.1365-2699.2008.02060.x>

- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R.; & Group, 1000 Genomes Project Analysis. (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>
- Denny, M. W., & Dowd, W. W. (2012). Biophysics, environmental stochasticity, and the evolution of thermal safety margins in intertidal limpets. *Journal of Experimental Biology*, 215(6), 934–947. <https://doi.org/10.1242/jeb.058958>
- Detrich, H. W., Parker, S. K., Williams, J., Nogales, E., & Downing, K. H. (2000). Cold adaptation of microtubule assembly and dynamics. Structural interpretation of primary sequence changes present in the  $\alpha$ - and  $\beta$ -tubulins of antarctic fishes. *Journal of Biological Chemistry*, 275(47), 37038–37047. <https://doi.org/10.1074/jbc.M005699200>
- Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences of the United States of America*, 105(18), 6668–6672. <https://doi.org/10.1073/pnas.0709472105>
- Dillon, M. E., Wang, G., Garrity, P. A., & Huey, R. B. (2009). Thermal preference in *Drosophila*. *Journal of Thermal Biology*, 34(3), 109–119. <https://doi.org/10.1016/j.jtherbio.2008.11.007>
- Dittmar, E. L., Oakley, C. G., Conner, J. K., Gould, B. A., & Schemske, D. W. (2016). Factors influencing the effect size distribution of adaptive substitutions. *Proceedings of the Royal Society B: Biological Sciences*, 283(1828), 20153065. <https://doi.org/10.1098/rspb.2015.3065>
- Divino, J. N., Monette, M. Y., McCormick, S. D., Yancey, P. H., Flannery, K. G., Bell, M. A., Rollins, J. L., Von Hippel, F. A., & Schultz, E. T. (2016). Osmoregulatory physiology and rapid evolution of salinity tolerance in threespine stickleback recently introduced to fresh water. *Evolutionary Ecology Research*, 17(2), 179–201.
- Eliason, E. J., Clark, T. D., Hague, M. J., Hanson, L. M., Gallagher, Z. S., Jeffries, K. M., Gale, M. K., Patterson, D. A., Hinch, S. G., & Farrell, A. P. (2011). Differences in thermal tolerance among sockeye salmon populations. *Science*, 332(6025), 109–112. <https://doi.org/10.1126/science.1199158>
- Environmental Systems Research Institute. (2017). ArcGIS Desktop. Release, 10(8).
- Etterson, J. R., & Shaw, R. G. (2001). Constraint to adaptive evolution in response to global warming. *Science*, 294(5540), 151–154. <https://doi.org/10.1126/science.1063656>
- Evans, T. G., Diamond, S. E., & Kelly, M. W. (2015). Mechanistic species distribution modelling as a link between physiology and conservation. *Conservation Physiology*, 3(1), 1–16. <https://doi.org/10.1093/conphys/cov056>
- Everett, M. V., & Seeb, J. E. (2014). Detection and mapping of QTL for temperature tolerance and body size in Chinook salmon (*Oncorhynchus tshawytscha*) using genotyping by sequencing. *Evolutionary Applications*, 7(4), 480–492. <https://doi.org/10.1111/eva.12147>
- Eyer, P. A., Blumenfeld, A. J., & Vargo, E. L. (2019). Sexually antagonistic selection promotes genetic divergence between males and females in an ant. *Proceedings of the National Academy of Sciences of the United States of America of the United States of America*, 116(48), 24157–24163. <https://doi.org/10.1073/pnas.1906568116>
- Fangue, N. A., Hofmeister, M., & Schulte, P. M. (2006). Intraspecific variation in thermal tolerance and heat shock protein gene expression in common killifish, *Fundulus heteroclitus*. *The Journal of Experimental Biology*, 209(15), 2859–2872. <https://doi.org/10.1242/jeb.02260>
- Fangue, N. A., Richards, J. G., & Schulte, P. M. (2009). Do mitochondrial properties explain intraspecific variation in thermal tolerance? *The Journal of Experimental Biology*, 212(4), 514–522. <https://doi.org/10.1242/jeb.024034>
- Fetterer, F., Savoie, M., Helfrich, S., & Clemene-Colon, P. (2010). U.S. national ice center and national snow and ice data center. In *Multisensor analyzed sea ice extent - northern hemisphere (MASIE-NH)*. National Snow and Ice Data Center. <https://doi.org/10.7265/N5GT5K3K>
- Fisher, R. (1930). *The genetical theory of natural selection*. 1st edn. The Clarendon Press.
- Fossheim, M., Primicerio, R., Johannesen, E., Ingvaldsen, R. B., Aschan, M. M., & Dolgov, A. V. (2015). Recent warming leads to a rapid borealization of fish communities in the Arctic. *Nature Climate Change*, 5(7), 673–677. <https://doi.org/10.1038/nclimate2647>
- Gilchrist, G. W., & Huey, R. B. (1999). The direct response of *Drosophila melanogaster* to selection on knockdown temperature. *Heredity*, 83(1), 15–29. <https://doi.org/10.1038/sj.hdy.6885330>
- Gómez-Ruiz, E. P., & Lacher, T. E. (2019). Climate change, range shifts, and the disruption of a pollinator-plant complex. *Scientific Reports*, 9(1), 1–10. <https://doi.org/10.1038/s41598-019-53670-9>
- Greenwood, A. K., Jones, F. C., Chan, Y. F., Brady, S. D., Absher, D. M., Grimwood, J., Schmutz, J., Myers, R. M., Kingsley, D. M., & Peichel, C. L. (2011). The genetic basis of divergent pigment patterns in juvenile threespine sticklebacks. *Heredity*, 107(2), 155–166. <https://doi.org/10.1038/hdy.2011.1>
- Griswold, C. K., & Whitlock, M. C. (2003). The genetics of adaptation: The roles of pleiotropy, stabilizing selection and drift in shaping the distribution of bidirectional fixed mutational effects. *Genetics*, 165(4), 2181–2192. <https://doi.org/10.1093/genetics/165.4.2181>
- Guisan, A., Tingley, R., Baumgartner, J. B., Naujokaitis-Lewis, I., Sutcliffe, P. R., Tulloch, A. I. T., Regan, T. J., Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T. G., Rhodes, J. R., Maggini, R., Setterfield, S. A., Elith, J., Schwartz, M. W., Wintle, B. A., Broennimann, O., Austin, M., ... Buckley, Y. M. (2013). Predicting species distributions for conservation decisions. *Ecology Letters*, 16(12), 1424–1435. <https://doi.org/10.1111/ele.12189>
- Hangartner, S., & Hoffmann, A. A. (2016). Evolutionary potential of multiple measures of upper thermal tolerance in *Drosophila melanogaster*. *Functional Ecology*, 30(3), 442–452. <https://doi.org/10.1111/1365-2435.12499>
- Harrell, F. E. (2014). Hmisc: Harrell miscellaneous. R package v 3.14-0. <https://CRAN.R-project.org/package=Hmisc>
- Healy, T. M., Brennan, R. S., Whitehead, A., & Schulte, P. M. (2018). Tolerance traits related to climate change resilience are independent and polygenic. *Global Change Biology*, 24(11), 5348–5360. <https://doi.org/10.1111/gcb.14386>
- Hendry, A. P., Peichel, C. L., Matthews, B., Boughman, J. W., & Nosil, P. (2013). Stickleback research: The now and the next. *Evolutionary Ecology Research*, 15(2), 111–141.
- Herring, S., Christidis, N., Hoell, A., Hoerling, M., & Stott, P. (2020). Explaining extreme events of 2018 from a climate perspective. *Bulletin of the American Meteorological Society*, 101(1), 1–146. <https://doi.org/10.1175/1520-0477-95.9.s1.1>
- Herring, S., Christidis, N., Hoell, A., Kossin, J., Schreck, C., & Stott, P. (2018). Explaining extreme events of 2016 from a climate perspective. *Bulletin of the American Meteorological Society*, 99(1), S1–S157. <https://doi.org/10.1175/BAMS-ExplainingExtremeEvents2016.1>
- Herring, S., Hoerling, M., Kossin, J., Peterson, T., & Stott, P. (2015). Explaining extreme events of 2014 from a climate perspective. *Bulletin of the American Meteorological Society*, 96(12), 1–180. <https://doi.org/10.1175/BAMS-ExplainingExtremeEvents2014.1>
- Hijmans, R. J., van Etten, J., Sumner, M., Cheng, J., Bevan, A., Bevan, R., Busetto, L., Canty, M., Forrest, D., Ghosh, A., Golicher, D., Gray, J., & Greenberg, J. A. (2020). raster: Geographic data analysis and modeling. <https://cran.r-project.org/web/packages/raster/raster.pdf>
- Hochachka, P. W., & Somero, G. N. (2002). Mechanism and process in physiological evolution. *Biochemical Adaptation*, 480.
- Hoffman, A., & Sgrò, C. (2011). Climate change and evolutionary adaptation. *Nature*, 470, 479–485. <https://doi.org/10.1038/nature09670>

- Horton, K. G., La Sorte, F. A., Sheldon, D., Lin, T. Y., Winner, K., Bernstein, G., Maji, S., Hochachka, W. M., & Farnsworth, A. (2020). Phenology of nocturnal avian migration has shifted at the continental scale. *Nature Climate Change*, 10(1), 63–68. <https://doi.org/10.1038/s41558-019-0648-9>
- Hovel, R. A., Carlson, S. M., & Quinn, T. P. (2016). Climate change alters the reproductive phenology and investment of a lacustrine fish, the three-spine stickleback. *Global Change Biology*, 23(6), 2308–2320. <https://doi.org/10.1111/gcb.13531>
- Hu, D., Guan, Z., Tian, W., & Ren, R. (2018). Recent strengthening of the stratospheric Arctic vortex response to warming in the central North Pacific. *Nature Communications*, 9(1). <https://doi.org/10.1038/s41467-018-04138-3>
- Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., & Williams, S. E. (2012). Predicting organismal vulnerability to climate warming: Roles of behaviour, physiology and adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 1665–1679. <https://doi.org/10.1098/rstb.2012.0005>
- Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm performance. *Trends in Ecology & Evolution*, 4(5), 131–135. [https://doi.org/10.1016/0169-5347\(89\)90211-5](https://doi.org/10.1016/0169-5347(89)90211-5)
- Huey, R. B., & Kingsolver, J. G. (2011). Variation in universal temperature dependence of biological rates. *Proceedings of the National Academy of Sciences of the United States of America*, 108(26), 10377–10378. <https://doi.org/10.1073/pnas.1107430108>
- Huey, R. B., & Slatkin, M. (1976). Costs and benefits of lizard thermoregulation. *Quarterly Review of Biology*, 51(3), 363–384.
- Hutchison, V. H. (1961). Comparative biology critical thermal maxima in salamanders. *Physiological Zoology*, 34(2), 92–125. <https://doi.org/10.1086/physzool.34.2.30152688>
- IPCC. (2014). Climate change 2014: Synthesis report. In *Fifth assessment report of the Intergovernmental Panel on Climate Change*. [https://doi.org/10.1016/S0022-0248\(00\)00575-3](https://doi.org/10.1016/S0022-0248(00)00575-3)
- IPCC. (2018). *IPCC special report on the impacts of global warming of 1.5°C - Summary for policy makers*. <http://www.ipcc.ch/report/sr15/>
- Jackson, T. R., Ferguson, M. M., Danzmann, R. G., Fishback, A. G., Ihssen, P. E., O'Connell, M., & Crease, T. J. (1998). Identification of two QTL influencing upper temperature tolerance in three rainbow trout (*Oncorhynchus mykiss*) half-sib families. *Heredity*, 80(2), 143–151. <https://doi.org/10.1038/sj.hdy.6882890>
- Jin, Y., Zhou, T., Geng, X., Liu, S., Chen, A., Yao, J., Jiang, C., Tan, S., Su, B., & Liu, Z. (2017). A genome-wide association study of heat stress-associated SNPs in catfish. *Animal Genetics*, 48(2), 233–236. <https://doi.org/10.1111/age.12482>
- Johannessen, O. M., Bengtsson, L., Miles, M. W., Kuzmina, S. I., Semenov, V. A., Alekseev, G. V., Nagurnyi, A. P., Zakharov, V. F., Bobylev, L. P., Pettersson, L. H., Hasselmann, K., & Cattle, H. P. (2004). Arctic climate change: observed and modelled temperature and sea-ice variability. *Tellus A: Dynamic Meteorology and Oceanography*, 56(4), 328–341. <https://doi.org/10.3402/tellusa.v56i4.14418>
- Jones, F. C., Chan, Y. F., Schmutz, J., Grimwood, J., Brady, S. D., Southwick, A. M., Absher, D. M., Myers, R. M., Reimchen, T. E., Deagle, B. E., Schluter, D., & Kingsley, D. M. (2012). A genome-wide SNP genotyping array reveals patterns of global and repeated species-pair divergence in sticklebacks. *Current Biology*, 22(1), 83–90. <https://doi.org/10.1016/j.cub.2011.11.045>
- Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Muceli, E., Johnson, J., Swofford, R., Pirun, M., Zody, M. C., White, S., Birney, E., Searle, S., Schmutz, J., Grimwood, J., Dickson, M. C., Myers, R. M., Miller, C. T., Summers, B. R., Knecht, A. K., ... Kingsley, D. M. (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, 484(7392), 55–61. <https://doi.org/10.1038/nature10944>
- Kearney, M., & Porter, W. P. (2004). Mapping the fundamental niche: Physiology, climate, and the distribution of a nocturnal lizard. *Ecology*, 85(11), 3119–3131. <https://doi.org/10.1890/03-0820>
- Kearney, M., & Porter, W. (2009). Mechanistic niche modelling: Combining physiological and spatial data to predict species' ranges. *Ecology Letters*, 12(4), 334–350. <https://doi.org/10.1111/j.1461-0248.2008.01277.x>
- Kingsolver, J. G., Arthur Woods, H., Buckley, L. B., Potter, K. A., MacLean, H. J., & Higgins, J. K. (2011). Complex life cycles and the responses of insects to climate change. *Integrative and Comparative Biology*, 51(5), 719–732. <https://doi.org/10.1093/icb/1093/1093>
- Kirch, M., Romundset, A., Gilbert, M. T. P., Jones, F. C., & Foote, A. D. (2021). Ancient and modern stickleback genomes reveal the demographic constraints on adaptation. *Current Biology*, 31(9), 2027–2036.e8. <https://doi.org/10.1016/j.cub.2021.02.027>
- Kurz, W. A., Dymond, C. C., Stinson, G., Rampley, G. J., Neilson, E. T., Carroll, A. L., Ebata, T., & Safranyik, L. (2008). Mountain pine beetle and forest carbon feedback to climate change. *Nature*, 452(7190), 987–990. <https://doi.org/10.1038/nature06777>
- Larson, W. A., McKinney, G. J., Limborg, M. T., Everett, M. V., Seeb, L. W., & Seeb, J. E. (2016). Identification of multiple QTL hotspots in sockeye salmon (*Oncorhynchus nerka*) using genotyping-by-sequencing and a dense linkage map. *Journal of Heredity*, 107(2), 122–133. <https://doi.org/10.1093/jhered/escv099>
- Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics*, 26(5), 589–595. <https://doi.org/10.1093/bioinformatics/btp698>
- Lyon, N. J., Debinski, D. M., & Rangwala, I. (2019). Evaluating the utility of species distribution models in informing climate change-resilient grassland restoration strategy. *Frontiers in Ecology and Evolution*, 7(February), 1–8. <https://doi.org/10.3389/fevo.2019.00033>
- McLeod, D. J., Hallegraeff, G. M., Hosie, G. W., & Richardson, A. J. (2012). Climate-driven range expansion of the red-tide dinoflagellate *Noctiluca scintillans* into the Southern Ocean. *Journal of Plankton Research*, 34(4), 332–337. <https://doi.org/10.1093/plankt/fbr112>
- Melles, S. J., Fortin, M. J., Lindsay, K., & Badzinski, D. (2011). Expanding northward: Influence of climate change, forest connectivity, and population processes on a threatened species' range shift. *Global Change Biology*, 17(1), 17–31. <https://doi.org/10.1111/j.1365-2486.2010.02214.x>
- Morgan, R., Finnøen, M. H., Jensen, H., Pélabon, C., & Jutfelt, F. (2020). Low potential for evolutionary rescue from climate change in a tropical fish. *Proceedings of the National Academy of Sciences of the United States of America*, 117(52), 33365–33372. <https://doi.org/10.1073/PNAS.2011419117>
- Morgan, R., Finnøen, M. H., & Jutfelt, F. (2018). CTmax is repeatable and doesn't reduce growth in zebrafish. *Scientific Reports*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-25593-4>
- Morgan, R., Sundin, J., Finnøen, M. H., Dresler, G., Martínex Vendrell, M., Dey, A., Sarkar, K., & Jutfelt, F. (2019). Are model organisms representative for climate change research? Testing thermal tolerance in wild and laboratory zebrafish populations. *Conservation Physiology*, 7(1), coz036. <https://doi.org/10.1093/conphys/coz036>
- Morris, M. R. J., Bowles, E., Allen, B. E., Jamniczky, H. A., & Rogers, S. M. (2018). Contemporary ancestor? Adaptive divergence from standing genetic variation in Pacific marine threespine stickleback. *BMC Evolutionary Biology*, 18(1), 1–21. <https://doi.org/10.1186/s12862-018-1228-8>
- Moyano, M., Candebat, C., Ruhbaum, Y., Álvarez-Fernández, S., Claireaux, G., Zambonino-Infante, J. L., & Peck, M. A. (2017). Effects of warming rate, acclimation temperature and ontogeny on the critical thermal maximum of temperate marine fish larvae. *PLoS One*, 12(7), 1–23. <https://doi.org/10.1371/journal.pone.0179928>
- Muñoz, N. J., Farrell, A. P., Heath, J. W., & Neff, B. D. (2014). Adaptive potential of a Pacific salmon challenged by climate change. *Nature Climate Change*, 5(2), 163–166. <https://doi.org/10.1038/nclimate2473>

- Nati, J. J. H., Svendsen, M. B. S., Marras, S., Killen, S. S., Steffensen, J. F., McKenzie, D. J., & Domenici, P. (2021). Intraspecific variation in thermal tolerance differs between tropical and temperate fishes. *Scientific Reports*, 11(1), 1–8. <https://doi.org/10.1038/s41598-021-00695-8>
- Ogden, N. H., Maarouf, A., Barker, I. K., Bigras-Poulin, M., Lindsay, L. R., Morshed, M. G., O'Callaghan, C. J., Ramay, F., Waltner-Toews, D., & Charron, D. F. (2006). Climate change and the potential for range expansion of the Lyme disease vector *Ixodes scapularis* in Canada. *International Journal for Parasitology*, 36(1), 63–70. <https://doi.org/10.1016/j.ijpara.2005.08.016>
- Orr, H. A. (1998). The population genetics of adaptation: The distribution of factors fixed during adaptive evolution. *Evolution*, 52(4), 935–949. <https://doi.org/10.1111/j.1558-5646.1998.tb01823.x>
- Orr, H. A. (2005). The genetic theory of adaptation: A brief history. *Nature Reviews Genetics*, 6(2), 119–127. <https://doi.org/10.1038/nrg1523>
- Ouellette, L. A., Reid, R. W., Blanchard, S. G., & Brouwer, C. R. (2018). LinkageMapView-rendering high-resolution linkage and QTL maps. *Bioinformatics*, 34(2), 306–307. <https://doi.org/10.1093/bioinformatics/btx576>
- Parnesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421, 37–42. <https://doi.org/10.1038/nature01286>
- Perry, G. M. L., Danzmann, R. G., Ferguson, M. M., & Gibson, J. P. (2001). Quantitative trait loci for upper thermal tolerance in outbred strains of rainbow trout (*Oncorhynchus mykiss*). *Heredity*, 86(3), 333–341. <https://doi.org/10.1046/j.1365-2540.2001.00838.x>
- Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One*, 7(5). <https://doi.org/10.1371/journal.pone.0037135>
- Platts, P. J., Mason, S. C., Palmer, G., Hill, J. K., Oliver, T. H., Powney, G. D., Fox, R., & Thomas, C. D. (2019). Habitat availability explains variation in climate-driven range shifts across multiple taxonomic groups. *Scientific Reports*, 9(1), 1–10. <https://doi.org/10.1038/s41598-019-51582-2>
- Puritz, J. B., Hollenbeck, C. M., & Gold, J. R. (2014). *dDocent*: A RADseq, variant-calling pipeline designed for population genomics of non-model organisms. *PeerJ*, 2, e431. <https://doi.org/10.7717/peerj.431>
- R Core Team. (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rastas, P. (2017). Lep-MAP3: Robust linkage mapping even for low-coverage whole genome sequencing data. *Bioinformatics*, 33(23), 3726–3732. <https://doi.org/10.1093/bioinformatics/btx494>
- Reynolds, R., Smith, T., Liu, C., Chelton, D., Casey, K., & Schlax, M. (2007). Daily high-resolution-blended analyses for sea surface temperature. *Journal of Climate*, 20(22), 5473–5496. <https://doi.org/10.1175/2007jcli1824.1>
- Roger, A., Stuetz, R., Ove, K., Giraudoux, P., & Santilli, S. (2020). *rgeos: Interface to geometry engine—Open*.
- Rogers, S. M., Tamkee, P., Summers, B., Balabhadra, S., Marks, M., Kingsley, D. M., & Schluter, D. (2012). Genetic signature of adaptive peak shift in threespine stickleback. *Evolution*, 66(12), 2439–2451. <https://doi.org/10.5061/dryad.6jj614kh>
- Rolandi, C., Lighton, J. R. B., de la Vega, G. J., Schilman, P. E., & Mensch, J. (2018). Genetic variation for tolerance to high temperatures in a population of *Drosophila melanogaster*. *Ecology and Evolution*, 8(21), 10374–10383. <https://doi.org/10.1002/ece3.4409>
- Rushing, C. S., Andrew Royle, J., Ziolkowski, D. J., & Pardieck, K. L. (2020). Migratory behavior and winter geography drive differential range shifts of eastern birds in response to recent climate change. *Proceedings of the National Academy of Sciences of the United States of America*, 117(23), 12897–12903. <https://doi.org/10.1073/pnas.2000299117>
- Sagurova, I., Ludwig, A., Ogden, N. H., Pelcat, Y., Dueymes, G., & Gachon, P. (2019). Predicted northward expansion of the geographic range of the tick vector *Amblyomma americanum* in North America under future climate conditions. *Environmental Health Perspectives*, 127(10), 1–14. <https://doi.org/10.1289/EHP5668>
- Sambaraju, K. R., Carroll, A. L., & Aukema, B. H. (2019). Multiyear weather anomalies associated with range shifts by the mountain pine beetle preceding large epidemics. *Forest Ecology and Management*, 438, 86–95. <https://doi.org/10.1016/j.foreco.2019.02.011>
- Sanderson, S., Beausoleil, M., O'Dea, R. E., Wood, Z. T., Correa, C., Frankel, V., Gorné, L. D., Haines, G. E., Kinnison, M. T., Oke, K. B., Pelletier, F., Pérez-Jvostov, F., Reyescorral, W. D., Ritchot, Y., Sorbara, F., Gotanda, K. M., & Hendry, A. P. (2021). The pace of modern life, revisited. *Molecular Ecology*, 31(4), 1028–1043. <https://doi.org/10.1111/mec.16299>
- Schluter, D., & Conte, G. L. (2009). Genetics and ecological speciation. *Proceedings of the National Academy of Sciences of the United States of America*, 3, 47–64. <https://doi.org/10.17226/12692>
- Schluter, D., Marchinko, K. B., Arnegard, M. E., Zhang, H., Brady, S. D., Jones, F. C., Bell, M. A., & Kingsley, D. M. (2021). Fitness maps to a large-effect locus in introduced stickleback populations. *Proceedings of the National Academy of Sciences of the United States of America*, 118(3), 1–10. <https://doi.org/10.1073/pnas.1914889118>
- Schluter, D., Marchinko, K., Barrett, R., & Rogers, S. (2010). Natural selection and the genetics of adaptation in threespine stickleback. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1552), 2479–2486. <https://doi.org/10.1098/rstb.2010.0036>
- Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and species range limits. *Annual Review of Ecology, Evolution, and Systematics*, 40(1), 415–436. <https://doi.org/10.1146/annurev.ecolsys.110308.120317>
- Shin, S. C., Ahn, D. H., Kim, S. J., Pyo, C. W., Lee, H., Kim, M.-K., Lee, J., Lee, J. E., Detrich, H. W., Postlethwait, J. H., Edwards, D., Lee, S. G., Lee, J. H., & Park, H. (2014). The genome sequence of the Antarctic bullhead notothen reveals evolutionary adaptations to a cold environment. *Genome Biology*, 15(9), 468. <https://doi.org/10.1186/s13059-014-0468-1>
- Spies, I., Gruenthal, K. M., Drinan, D. P., Hollowed, A. B., Stevenson, D. E., Tarpey, C. M., & Hauser, L. (2020). Genetic evidence of a northward range expansion in the eastern Bering Sea stock of Pacific cod. *Evolutionary Applications*, 13(2), 362–375. <https://doi.org/10.1111/eva.12874>
- Stott, P. (2016). How climate change affects extreme weather events. *Science*, 352(6293), 1517–1518. <https://doi.org/10.1126/science.aaf7271>
- Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, 278(1713), 1823–1830. <https://doi.org/10.1098/rspb.2010.1295>
- Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, 2(9), 686–690. <https://doi.org/10.1038/nclimate1539>
- Tingley, M. W., Monahan, W. B., Beissinger, S. R., & Moritz, C. (2009). Birds track their Grinnellian niche through a century of climate change. *Proceedings of the National Academy of Sciences of the United States of America*, 106(Suppl. 2), 19637–19643. <https://doi.org/10.1073/pnas.0901562106>
- Tombre, I. M., Oudman, T., Shimmings, P., Griffin, L., & Prop, J. (2019). Northward range expansion in spring-staging barnacle geese is a response to climate change and population growth, mediated by individual experience. *Global Change Biology*, 25(11), 3680–3693. <https://doi.org/10.1111/gcb.14793>
- Turko, A. J., Nolan, C. B., Balshine, S., Scott, G. R., & Pitcher, T. E. (2020). Thermal tolerance depends on season, age and body condition

- in imperilled redbreasted dace *Clinostomus elongatus*. *Conservation Physiology*, 8(1), 1–15. <https://doi.org/10.1093/conphys/coaa062>
- Ungerer, M. C., & Riesebero, L. H. (2003). Genetic architecture of a selection response in *Arabidopsis thaliana*. *Evolution*, 57(11), 2531–2539. <https://doi.org/10.1111/j.0014-3820.2003.tb01497.x>
- Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M., Cornwell, W., Gianoli, E., van Kleunen, M., Naya, D. E., Nicotra, A. B., Poorter, H., & Zavala, M. A. (2014). The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecology Letters*, 17(11), 1351–1364. <https://doi.org/10.1111/ele.12348>
- Via, S., Conte, G., Mason-Foley, C., & Mills, K. (2012). Localizing FST outliers on a QTL map reveals evidence for large genomic regions of reduced gene exchange during speciation-with-gene-flow. *Molecular Ecology*, 21(22), 5546–5560. <https://doi.org/10.1111/mec.12021>
- Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J. M., Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, 416(6879), 389–395. <https://doi.org/10.1038/416389a>
- Weatherall, P., Marks, K., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J., Rovere, M., Chayes, D., Ferrini, V., & Wigley, R. (2015). A new digital bathymetric model of the world's oceans. *Earth and Space Science*, 2(8), 331–345. <https://doi.org/10.1002/2015EA000107>
- Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., & Zemla, J. (2017). Package 'corrplot'. *Statistician*, 56, 316–324.
- Wiens, J., Stralberg, D., Jongsomjit, D., Howell, C., & Snyder, M. (2009). Niches, models, and climate change: Assessing the assumptions and uncertainties. *Proceedings of the National Academy of Sciences of the United States of America*, 106(Supplement\_2), 19729–19736. <https://doi.org/10.1073/pnas.0901639106>
- Yapıcı, S., Bilge, G., & Filiz, H. (2016). Northwards range expansion of *Sparisoma cretense* (Linnaeus, 1758) in the Turkish Aegean Sea. *Journal of Aquaculture Engineering and Fisheries Research*, 201–207. <https://doi.org/10.3153/jaefr16022>
- Yeaman, S., & Otto, S. P. (2011). Establishment and maintenance of adaptive genetic divergence under migration, selection, and drift. *Evolution*, 65(7), 2123–2129. <https://doi.org/10.1111/j.1558-5646.2011.01277.x>
- Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-selection balance. *Evolution*, 65(7), 1897–1911. <https://doi.org/10.1111/j.1558-5646.2011.01269.x>
- Zeidberg, L. D., & Robison, B. H. (2007). Invasive range expansion by the Humboldt squid, *Dosidicus gigas*, in the eastern North Pacific. *Proceedings of the National Academy of Sciences of the United States of America*, 104(31), 12948–12950. <https://doi.org/10.1073/pnas.0702043104>
- Zweng, M. M., Reagan, J. R., Antonov, J. I., Mishonov, A. V., Boyer, T. P., Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., & Bidlle, M. M. (2013). World ocean atlas 2013, volume 2: Salinity. In NOAA Atlas NESDIS 74 (Vol. 2, Issue 1). <https://doi.org/10.1182/blood-2011-06-357442>

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Smith, S. J., Mogensen, S., Barry, T. N., Paccard, A., Jamniczky, H. A., Barrett, R. D. H., & Rogers, S. M. (2022). Evolution of thermal physiology alters the projected range of threespine stickleback under climate change. *Molecular Ecology*, 31, 2312–2326. <https://doi.org/10.1111/mec.16396>